

ARTICLES

Endotoxin and immune activation in chronic heart failure: a prospective cohort study

Josef Niebauer, Hans-Dieter Volk, Michael Kemp, Martin Dominguez, Ralf R Schumann, Mathias Rauchhaus, Philip A Poole-Wilson, Andrew J S Coats, Stefan D Anker

Summary

Background Immune activation in patients with chronic heart failure may be secondary to endotoxin (lipopolysaccharide) action. We investigated the hypothesis that altered gut permeability with bacterial translocation and endotoxaemia would be increased in patients with oedema secondary to congestive heart failure.

Methods We compared 20 patients who had chronic heart failure with recent-onset peripheral oedema (mean age 64 years [SD 10], New York Heart Association [NYHA] class 3-3 [0-7]), 20 stable non-oedematous patients with chronic heart failure (mean age 63 years [19], NYHA class 2-6 [0-7]), and 14 healthy volunteers (mean age 55 years [16]). Biochemical markers of endotoxaemia, inflammation, and immune activation were measured. Ten patients were studied within 1 week of complete resolution of oedema. Five patients survived longer than 6 months and were restudied again after remaining free of oedema for more than 3 months.

Findings Mean endotoxin concentrations were higher in oedematous patients with chronic heart failure than in stable patients with chronic heart failure (0.74 [SD 0.45] vs 0.37 EU/mL [0.23], $p=0.009$) and controls (0.46 EU/mL [0.21], $p=0.02$). Oedematous patients had the highest concentrations of several cytokines. After short-term diuretic treatment, endotoxin concentrations decreased from 0.84 EU/mL [0.49] to 0.45 EU/mL [0.21], $p<0.05$ but cytokines remained raised. After freedom of oedema for more than 3 months after oedema resolved, endotoxin concentrations remained unchanged from the previous visit (0.49 EU/mL [0.06], $p=0.45$).

Interpretation Raised concentrations of endotoxin and cytokines are found in patients with chronic heart failure during acute oedematous exacerbation. Intensified diuretic treatment can normalise endotoxin concentrations. Our preliminary findings suggest that endotoxin may trigger immune activation in patients with chronic heart failure during oedematous episodes.

Lancet 1999; 353: 1838-42

See Commentary page 1812

Cardiac Medicine, National Heart and Lung Institute, Imperial College School of Medicine, London, UK (J Niebauer MD, M Kemp MD, M Rauchhaus, Prof P A Poole-Wilson MD, Prof A J S Coats MD, S D Anker MD); Karlsruher Herzszentrum der Universität Leipzig, Germany (J Niebauer); Institut für Medizinische Immunologie, Universitätsklinikum Charité, Berlin, Germany (Prof HD Volk MD); Heart Science Center, Harfield Hospital, UK (M Dominguez MD); Institut für Mikrobiologie und Hygiene, Universitätsklinikum Charité, Berlin (R R Schumann MD); and Fraunhofer-Klinik am Max-Deebüch Zentrum für Molekulare Medizin, Charité, Campus Berlin-Buch, Berlin (S D Anker)

Correspondence to: Dr Stefan Anker, Cardiac Medicine, National Heart and Lung Institute, London SW3 6LY, UK (e-mail: s.anker@ic.nhs.uk).

Introduction

Some patients with chronic heart failure have features such as cardiac cachexia that may be due to activation of the immune system.^{1,2} Increased expression of tumour necrosis factor α (TNF α) has been found in cardiac tissue of patients with chronic heart failure undergoing heart transplantation and the failing heart has been suggested as the cause of immune activation.³ No link between a pathogenic process and cytokine activation has been documented in human beings with heart failure or in animal models. The cause of increased cytokine production in patients with heart failure remains unknown.

We have previously suggested that bacterial endotoxin, lipopolysaccharide, contributes to immune activation in chronic heart failure.⁴ Acute venous congestion could lead to altered gut permeability for bacteria, endotoxin, or both, and to translocation of these materials into the circulation. In the circulation, lipopolysaccharide is bound by a serum protein, termed lipopolysaccharide-binding protein (LBP).⁵ The lipopolysaccharide-LBP complex can interact with the CD14 membrane protein and Toll-like signalling receptors to start a signalling cascade that leads to increased cytokine production (figure 1). The extracellular domain of the CD14 receptor is shed after interaction and serum concentrations are thought to reflect the amount of endotoxin and cell interaction. The lipopolysaccharide-LBP ratio has been shown to be crucial for the immunostimulatory effects of lipopolysaccharide.⁶ High concentrations of LBP, as seen during the acute-phase response, can completely block lipopolysaccharide effects in vitro and in a murine sepsis model.⁷ Furthermore, patients with high concentrations of soluble CD14 (which shows endotoxin-cell interaction and shedding of CD14 from the cell membrane⁸) have strikingly increased concentrations of TNF α , soluble TNF receptor-1 and receptor-2, and intracellular-adhesion molecule-1.⁴

The degree of bowel-wall oedema cannot be directly measured. The relation between central haemodynamics and the pathophysiological features of chronic heart failure is weak.⁹ In animal models there is a poor relation between intracardiac pressures and intestinal perfusion.¹⁰ We therefore separated patients according to the presence or absence of a reliable marker of acute venous congestion due to cardiac failure, namely peripheral oedema. Bowel-wall oedema that could cause altered gut permeability and bacterial (ie, endotoxin) translocation is most likely to occur with moderate to severe peripheral oedema.

Our main aim in this study was to measure endotoxin and cytokine concentrations in patients with chronic heart failure during an acute exacerbation with peripheral oedema and after short-term and long-term treatment with diuretics.

Methods

Participants

We studied prospectively 14 healthy volunteers (mean age 55 years [SD 16]) and 40 patients with chronic heart failure (mean age 63 years [15], $p=0.11$). We did the baseline studies between April and October, 1997. 20 stable patients were recruited during outpatient clinics on 3 specific days and 20 patients with moderate

```

graph TD
    A[Congestive heart failure] --> B[Increased venous congestion  
(left heart failure) or right]
    B --> C[Lipopolysaccharide is translocated to  
peritoneal macrophages, if both]
    C --> D[Discompartments (peritoneal) effluent:  
Increased bacteria (LPS)  
concentrations in effluent]
    D --> E[LPS LBP clustering LBP binds to LPS]
    E --> F[CD14 (receptor) LPS-LBP complex Toll-like receptor  
Macrophage, fibrocyte, dendritic cells]
    F --> G[Signal transduction and  
subsequent cytokine production  
(i.e., tumor necrosis factor)]
    G --> H[Increased concentrations of  
TNF in effluent]
    F --> I[Exaggerated duration of CD14 receptor to  
cells (soluble CD14 can be recognized  
in effluent)]
  
```

After a median of 14 days (range 7-89) we restudied these patients within 1 week of complete resolution of oedema (after treatment six NYHA class III, four class II; mean weight loss 3.6 kg [0.3, range 2.5-5.0]). Nine patients regained clinical

In a subgroup of ten non-odentogenic and seven odontogenic patients, as well as in all healthy volunteers, whole-blood samples were taken in potassium citric-acid tubes (Vacutainer Systems, Falco HD, Oxford, UK) and stained with fluorescently labelled monoclonal antibodies (Coulter Electronics, Luton, UK) to

	Healthy volunteers (n=14)	CHP-10 patients (n=20)	CHP-10 controls (n=20)
Demographics (mean (SD))			
Age (years)	58 (18)	63 (15)	64 (15)
Weight (kg)	74 (7)	76 (8)	78 (8)
NYHA class	..	2-6 (3-7)	3-3 (3-3)
Drugs			
Ischemic	..	15	11
Hypertensive	..	4	9
Clinical (mean (SD))			
Sodium (mmol/L)	139 (3-5)	137 (4-5)	134 (4-5)
Creatinine (µmol/L)	82 (19)	131 (68)	219 (100)*
Urea (mmol/L)	5-6 (0-8)	11-6 (7-9)	20-0 (12-4)*
Uric acid (mmol/L)	308 (83)	417 (146)	640 (174)*
Aspartate aminotransferase (U/L)	28 (14)	24 (5)	23 (5)
Aspartate aminotransferase (U/L)	23 (10)	17 (4)	14 (4)
Lymphocyte profile (mean (SD))			
CD4	47 (5)	61 (11)	35 (15)*
CD8	22 (7)	23 (14)	28 (20)
CD4/8 ratio	2-3 (1-2)	3-3 (2-4)	2-6 (2-3)
CD4/25 ratio	6-7 (4-2)	5-5 (2-4)	10-6 (8-7)
CD8/25 ratio	4-7 (2-4)	8-7 (4-6)	21-6 (10-7)*

Table 1: Characteristics of patients with chronic heart failure with and without peripheral oedema and healthy volunteers

ARTICLES

	Healthy volunteers (n=14)	CHF no oedema (n=20)	CHF oedema (n=20)
Endotoxin (EU/mL)	0.46 (0.21)	0.37 (0.23)	0.74 (0.43)*
LBP (pg/mL)	8.8 (4.6)	10.4 (9.3)	12.1 (9.0)
Lipopolysaccharide/kg LBP ratio	0.64 (0.20)	0.44 (0.30)	0.75 (0.48)
TNF α (pg/mL)	24.8 (9.6)	35.8 (7.0)	35.6 (12.3)*
Soluble TNF receptor-1 (pg/mL)	708 (213)	1077 (529)	1822 (1398)*
Soluble TNF receptor-2 (pg/mL)	1485 (839)	2085 (1380)	3143 (1690)*
Soluble CD14 (ng/mL)	2459 (583)	3574 (454)	4243 (580)*
Procalcitonin (pg/mL)	87 (1.6)	208 (7.3)	345 (9.4)
Interleukin-6 (pg/mL)	2.0 (0.4)	4.3 (0.6)	14.7 (17.3)*
C-reactive protein (mg/L)	5.8 (1.7)	8.5 (0.5)	19.7 (17.1)*

* $p < 0.05$ vs healthy volunteers, † $p < 0.01$ vs CHF no oedema, ‡ $p < 0.01$ vs CHF no oedema, § $p < 0.01$ vs healthy volunteers, ¶ $p < 0.001$ vs healthy volunteers, * $p < 0.05$ vs CHF no oedema.

Table 2: Mean (SD) plasma concentrations of endotoxin and inflammatory markers in healthy volunteers and patients with chronic heart failure

determine peripheral lymphocyte phenotype and the proportion of CD25 receptor-positive T cells. A staining excess of antibody, determined by titration (data not shown), was placed into 12x75 mm polypropylene tubes (Bibby, Hampshire, UK). Two tubes were analysed for each patient's sample. The first tube contained control monoclonal mouse anti-human antibodies isotopically matched to the test antibodies in the second tube. The antibody-fluorescence conjugates used were CD3-PC5, CD4-FITC, CD8-BOD, and CD25R-RD1. The fume-acid lysed whole-blood protocol was used in the multi-Q-prep (Coulter Electronics, Luton, UK). Lymphocyte gating was set on forward compared with side-scatter dot plot, and compensation was established by the combining of single-colour-stained leucocyte populations. Four-colour flow cytometry was done on the Cocker XL-MCL with Sysmex II software (version 2.0).

Statistical analysis

We assessed normality of distribution with the Kolmogorov-Smirnov test. Unpaired Student's *t* test, paired *t* test, ANOVA with Fisher's post-hoc test (with allowance for multiple testing), and the Mann-Whitney *U* test were used where appropriate. Data are presented as mean (SD). We also used univariate-correlation and multivariate-correlation analyses to establish the relation between variables. We took $p < 0.05$ to be significant.

Results

In tables 1 and 2, baseline clinical characteristics and results of immunological and humoral measurements are shown. Endotoxin concentrations were highest in heart-failure patients with peripheral oedema, compared with heart-failure patients without oedema (98%) and controls (59%, $p = 0.0027$; figure 2). Plasma concentrations of LBP did not differ between groups, but there was a raised lipopolysaccharide/kg LBP ratio in the heart-failure patients with oedema compared with those without oedema (71%, $p < 0.01$). In oedematous heart-failure patients, plasma concentrations were significantly higher for C-reactive protein, TNF α , soluble TNF receptor-1 and receptor-2, interleukin-6, and soluble CD14 (table 2) than for all other groups.

Among all participants ($n = 54$), concentrations of soluble CD14 correlated significantly with endotoxin ($r = 0.30$, $p < 0.05$). This correlation was not significant when patients or healthy volunteers were analysed separately. In all patients with chronic heart failure, soluble CD14 correlated with TNF α ($r = 0.32$, $p < 0.05$) and soluble TNF receptor-1 ($r = 0.43$, $p < 0.01$). There was a correlation between soluble CD14 and soluble TNF receptor-2 in patients with stable chronic heart failure ($r = 0.61$, $p < 0.01$).




Figure 2: Plasma endotoxin concentrations in healthy volunteers and heart failure patients with and without oedema. Short horizontal lines=10th and 90th percentiles; long horizontal lines=25th, 50th, and 75th percentiles.

No simple correlations existed between creatinine or uric acid plasma concentrations and lipopolysaccharides at baseline, nor between changes in markers of kidney function over time compared with changes of lipopolysaccharide or cytokine concentrations over time (data not shown). Therefore, a bias because of latent abnormalities of kidney function seen in some oedematous patients is unlikely.

Intensive diuretic treatment for a mean of 23 days (8) in ten patients with chronic heart failure resulted in a mean weight decrease of 3.6 kg (range 2.5–5.0), and improvement in the functional NYHA class in nine of the ten patients. In eight of these, endotoxin plasma concentration was decreased from 0.96 EU/mL (0.47) to 0.45 EU/mL (0.24). In two patients with normal concentrations of endotoxin at baseline, concentrations after diuretic treatment were 9% and 36% higher than at baseline, but still in the normal range (< 0.5 EU/mL). In all ten patients the lipopolysaccharide concentrations fell from 0.84 EU/mL (0.49) to 0.45 EU/mL (0.21), $p = 0.049$;

	Baseline (n=10)	After diuretic treatment (n=10)	p
Endotoxin (EU/mL)	0.84 (0.49)	0.45 (0.21)	<0.05
LBP (pg/mL)	10.3 (2.7)	12.7 (7.6)	0.37
Lipopolysaccharide/kg LBP ratio	0.85 (0.44)	0.28 (0.67)	0.033
TNF α (pg/mL)	39.8 (13.2)	40.2 (13.0)	0.82
Soluble TNF receptor-1 (pg/mL)	2138 (1314)	2785 (1392)	0.09
Soluble TNF receptor-2 (pg/mL)	3731 (1185)	4029 (1477)	0.48
Soluble CD14 (ng/mL)	4474 (527)	4430 (784)	0.69
Procalcitonin (pg/mL)	153 (32)	219 (104)	0.17
Interleukin-6 (pg/mL)	19.4 (23.0)	18.3 (23.6)	0.80
C-reactive protein (mg/L)	19.6 (11.0)	20.0 (20.7)	0.90

Table 3: Mean (SD) plasma concentrations of endotoxin and inflammatory markers in oedematous patients before and after diuretic treatment

ARTICLES




Figure 3: Effect of intensified diuretic treatment on plasma endotoxin concentrations in ten patients with chronic heart failure who had peripheral oedema. Short horizontal lines=25th and 75th percentiles; long horizontal lines=25th, 50th, and 75th percentiles; whiskers extend to 1.5th and 1.5th percentiles.

figure 3). The effect of diuretic treatment on the endotoxin and inflammatory markers are shown in table 3. During extended follow-up, five patients were re-assessed when free of oedema for more than 3 months after 21 weeks (7). Endotoxin concentrations at the third visit did not differ from those at the second visit after a mean of 19 days (0.39 [0.22] vs 0.49 EU/mL [0.06], $p=0.45$), but TNF α concentrations were lower (39.6 [12.4] vs 31.0 pg/mL [5.7], $p=0.079$).

Discussion

We have shown that endotoxin concentrations and proinflammatory cytokines are raised in patients with heart failure who have peripheral oedema. Raised endotoxin concentrations were normalised by heightened diuretic treatment. The endotoxaemia in these patients was not associated with a strong acute-phase response that would have led to an increased hepatic LBP synthesis and subsequent blocking of lipopolysaccharide effects. These results lend credence to the hypothesis that bacterial endotoxin may be an important stimulus of immune activation in patients with chronic heart failure. This finding may open various options for treatment of patients with chronic heart failure that could be directed against bacteria in the bowel, the translocation process, and endotoxin itself, the binding sites of bacterial endotoxin on immune competent cells, or both.

The complex of endotoxin and endotoxin-binding protein activates monocytes and tissue macrophages via the CD14 and Toll-like receptor proteins,¹⁰ which stimulates the production of TNF α and other cytokines (figure 1). Previous studies suggested that increased soluble CD14 concentrations might be related to endotoxaemia.⁴ We

established that oedematous patients have the highest concentrations of soluble CD14 and lipopolysaccharide, but in homogeneous groups of patients there was no direct numeric relation between the two variables. Shed, and therefore soluble, CD14 receptors are thought to reflect the amount of endotoxin/cell interaction in the long term. By contrast, endotoxin has a short plasma half-life (10–30 min), which may explain why soluble CD14 concentrations are more closely related to cytokines than endotoxin concentrations.⁴

The concentrations of endotoxin in our study were well below those seen in septic shock.¹⁰ Patients with chronic heart failure had no signs of active infection, and the moderate increases in plasma endotoxin are in keeping with the hypothesis of a translocation process. Possibly, it is endotoxin itself rather than bacteria that translocates. Lipopolysaccharides at baseline did not correlate significantly with renal function (as estimated by creatinine and urea) although this finding cannot completely exclude an influence of renal function on cytokine clearance.

Although intensified diuretic therapy resulted in normalisation of endotoxin concentrations, treatment did not lead immediately to lowered cytokine plasma concentrations, which is in keeping with a previous study.¹² This effect may be due to a concentration effect, resulting from the loss of up to 5 kg body water or long-term activation of monocytes or macrophages after brief exposure to an endotoxin stimulus during a phase of clinical deterioration with increased venous congestion. Alternatively, the lack of cytokine decrease immediately after clinical improvement may be due to a change in monocyte or macrophage lipopolysaccharide sensitivity (ie, "normalised" endotoxin concentrations may still cause increased cytokine production). Indeed, such an increased cellular sensitivity to lipopolysaccharides has been documented in patients with chronic heart failure who had acute decompensation.¹¹ The previously documented raised TNF α concentrations in cardiac tissue of patients with end-stage chronic heart failure may also be due to cardiomyocytes or tissue monocytes releasing increased amounts of cytokines upon stimulation by lipopolysaccharides because of decompensation or hypersensitive cardiomyocytes. In cardiomyocytes of heart transplantation recipients (especially in patients with ischaemic chronic heart failure) increased baseline and lipopolysaccharide-stimulated TNF α production has been reported.¹² In our study, after a long phase of clinical stability, TNF α plasma concentrations showed a strong trend to decrease back to normal, and, therefore, the process of normalisation of cytokine secretion seems to be slow.

Tolerance of monocytes or macrophages to endotoxin can be induced in vivo and in vitro by endotoxin itself. Such an effect frequently occurs after severe injury.¹³ One important mediator of lipopolysaccharides hyposensitivity is interleukin-10.¹⁴ Compared with controls, we found interleukin-10 to be lower in stable patients with chronic heart failure.⁴ Increased cardio-wall stress and general tissue hypoxia (both via local free-radical generation and subsequent stimulation of the nuclear factor- κ B pathway) and hormonal catabolic and anabolic imbalance (especially in patients with muscle wasting¹⁵) may lead to immunological hyposensitivity. Endotoxin may, therefore, be an important stimulus for cytokine production in the heart and in the periphery even in the absence of oedema. In-vitro low concentrations of lipopolysaccharides have

ARTICLES

detrimental effects on cardiomyocytes. These effects are indirect through the release of other substances,²¹ but direct effects also have been seen.²² In vivo there may be a dynamic balance between heart function and immune activation in patients with chronic heart failure, and over time patients with frequent oedematous episodes may deteriorate because of the cardiodepressant and metabolic consequences of raised TNF α concentrations. Better control of oedema in chronic heart failure may therefore be beneficial.

In stable ambulatory patients with chronic heart failure, a significant excess concentration of cytokines from the heart could not be shown,²³ which suggests that cardiac production may not be the main source of the raised peripheral cytokine plasma concentrations. In support of the importance of peripheral hypoxia, measures of increased oxidative stress have been found to correlate with soluble TNF receptor-1 and receptor-2 concentrations.²⁴ We have shown that peak leg blood flow after ischaemia in clinically stable patients with chronic heart failure is inversely related to TNF α plasma concentrations. This effect may be due to a relation between hypoxia and TNF α production or toxic effects of TNF α on endothelial function.²⁵ Hypoxia may not be the most important cytokine trigger in chronic heart failure because of the cytokine profile. Raised interleukin-6 plasma concentrations can be attributed to peripheral hypoxic conditions,²⁶ which will occur in chronic heart failure, but there is no report that hypoxia leads to raised concentrations of TNF α , procalcitonin, or soluble TNF receptor-1 or receptor-2. Soluble CD14 receptors are, by contrast, characteristic of endotoxin action, but not of hypoxic disorders.²⁷

This study shows the presence of raised plasma endotoxin concentrations in patients with chronic heart failure and peripheral oedema. In the presence of unchanged concentrations of endotoxin binding protein, the raised endotoxin concentration reflects a potentially pathogenic situation that leads to cytokine induction. We show that normalisation of endotoxin concentrations can be achieved by intensified diuretic treatment. Bacterial endotoxin may be an important stimulus of immune activation in patients with chronic heart failure. Our studies are preliminary and further investigations are needed. Nevertheless, these findings may open various new options for treatment directed against bacteria in the bowel, the translocation process, and endotoxin itself, the binding sites of bacterial endotoxin on immune competent cells, or both.

Contributors

Susien Anker and Andrew Coats developed the endotoxin hypothesis and with Philip Poole-Wilson designed the study. Josef Niebauer and Susien Anker coordinated the study. Josef Niebauer did all clinical measurements with the help of Matthias Rautavaara. Michel Kemp measured endotoxin and TNF α . Martin Dünnebier did all lymphocyte analyses. Hans-Dieter Volk and Ralf Schumann advised on immunological issues and measured all other cytokines and LBP. Josef Niebauer and Susien Anker analysed the data and prepared the manuscript with the help of Philip Poole-Wilson and Andrew Coats. Hans-Dieter Volk and Ralf Schumann edited the manuscript.

Acknowledgments

JN and SDA were supported by the Ernst und Bertha Grunke Stiftung, Düsseldorf, Germany. SDA is supported with a postgraduate fellowship of the Max-Debilität Centrum für Molekulare Medizin, Berlin, Germany. MR is supported by the Deutsche Forschungsgemeinschaft. AJS is supported by the Vascular Research Trust Fund. The Department of Cardiac Medicine is supported by the British Heart Foundation.

References

- 1 Levine B, Kahnman J, Mayer L, Füll H, Packer M. Elevated circulating levels of tumor necrosis factor in severe chronic heart failure. *N Engl J Med* 1990; 323: 236-41.
- 2 Anker SD, Swan JW, Chua TP, et al. Humoral changes and endothelial dysfunction in chronic heart failure: the importance of cardiac cachexia. *Circulation* 1997; 96: 536-44.
- 3 Tanc-Anders G, Kapadia S, Lee J, et al. Tumor necrosis factor- α and tumor necrosis factor receptors in the failing human heart. *Circulation* 1996; 93: 704-11.
- 4 Anker SD, Rector K, Volk HD, Kox WJ, Poole-Wilson PA, Coats AJS. Elevated soluble CD14 receptor and acute cytokines in chronic heart failure. *Am J Cardiol* 1997; 79: 1426-30.
- 5 Schwann RR, Leung SR, Flagg GW, et al. Structure and function of lipopolysaccharide binding protein. *Science* 1990; 249: 429-31.
- 6 Tobin PS, Golden K, Ivanc N, Hirsch R, Weiss R. Lipopolysaccharide (LPS) binding protein BPI and LBP from different types of complexes with LPS. *J Biol Chem* 1997; 272: 18682-85.
- 7 Lamping N, Dettmer R, Schröder NWJ, et al. LPS-binding protein protects mice from septic shock caused by LPS or gram-negative bacteria. *J Clin Invest* 1998; 101: 2063-71.
- 8 Ziegler-Heckroth, Ulich R. CD14: cell surface receptor and differentiation marker. *Immunol Today* 1993; 14: 121-25.
- 9 Clark AL, Poole-Wilson PA, Coats AJ. Bacterial lipopolysaccharide in chronic heart failure: central role of the periphery. *J Am Coll Cardiol* 1996; 28: 1092-102.
- 10 Swann T, Wessely C, Högler M, et al. Early effects of catecholamine therapy on mucosal integrity, intestinal blood flow, and oxygen metabolism in porcine endotoxin shock. *Ann Surg* 1998; 228: 235-48.
- 11 Lamping N, Hesse A, Yu B, et al. Effect of site-directed mutagenesis of guanine residues (Arg 94, Lys 95, Lys 99) of lipopolysaccharide (LPS)-binding protein on binding and transfer of LPS and subsequent immune cell activation. *J Immunol* 1994; 157: 4648-55.
- 12 Andros M, Gendrel D, Caron E, Raymond J, Gullward J, Bohman C. High serum procalcitonin concentrations in patients with sepsis and infection. *Lancet* 1993; 341: 515-18.
- 13 Yang RH, Mack MR, Gray A, et al. Toll-like receptor-2 mediates lipopolysaccharide-induced cellular signalling. *Nature* 1998; 395: 284-88.
- 14 Gomez-Ibanez J, Salgado A, Morillo M, et al. L-arginine nitric oxide pathway in endotoxaemia and human septic shock. *Crit Care Med* 1995; 23: 233-38.
- 15 Vandenbroucke M, Kienrich E, Paulus WJ. Pro-inflammatory cytokines and endothelium-dependent vasodilation in the forearm: serial assessment in patients with congestive heart failure. *Eur Heart J* 1998; 19: 747-52.
- 16 Vanhulst S, Bress B, Sella-Selzer M, Grunbach JM, Krumm H, Fügita HR. Monocyte activation in congestive heart failure due to coronary artery disease and idiopathic dilated cardiomyopathy. *Am J Cardiol* 1998; 81: 237-44.
- 17 Wagner DR, McIntyre C, Sanders VJ, Feldman AM. Adenosine inhibits lipopolysaccharide-induced secretion of tumor necrosis factor- α in the failing human heart. *Circulation* 1998; 97: 521-24.
- 18 Keel M, Schorcherberger N, Stockhuber U, et al. Endotoxin tolerance after severe injury and its regulatory mechanisms. *J Trauma* 1996; 41: 430-37.
- 19 Rando P, Sykes U, Michel C, et al. Mechanisms of endotoxin desensitization: involvement of interferon 10 and transforming growth factor beta. *J Exp Med* 1995; 181: 1187-92.
- 20 Anker SD, Clark AL, Kemp M, et al. Tumor necrosis factor and amyloid metabolism in chronic heart failure: possible relation to muscle wasting. *J Am Coll Cardiol* 1997; 30: 997-1001.
- 21 Lew WF, Ryan J, Yasuda S. Lipopolysaccharide induces cell shrinkage in rabbit ventricular cardiac myocytes. *Am J Physiol* 1997; 272: H2889-93.
- 22 Star RQ, Luder AS, Phillips GC, Strassman CB, Abel FL. Direct action of endotoxin on cardiac muscle. *Shock* 1995; 3: 380-84.
- 23 Mörner MA, Johansson B, Anker H, Collander KS, Gilbert EM. Circulating concentrations of proinflammatory cytokines in mild or moderate heart failure secondary to ischemic or idiopathic dilated cardiomyopathy. *Am J Cardiol* 1996; 77: 723-27.
- 24 Kishi M, Gokhale A, Solt MJ, et al. Increased oxidative stress in patients with congestive heart failure. *J Am Coll Cardiol* 1998; 31: 1352-56.
- 25 Anker SD, Vohringer M, Rector K, et al. Tumor necrosis factor alpha as a predictor of peak leg blood flow in patients with chronic heart failure. *Circulation* 1998; 98: 199-203.
- 26 Yin SP, Thibault I, Finley D, et al. Induction of interleukin-6 (IL-6) by hypoxia in vascular cells. *J Biol Chem* 1995; 270: 11463-71.
- 27 Eggertsen JR, Björnsdottir I, Lund PK, Jøe GB, Owebo R, Kjøniksen LP. LPS-induced release of IL-1 beta, IL-6, IL-8, TNF-alpha and sCD14 in whole blood and PBMC from patients with high or low levels of HDL. *Immunol* 1994; 6: 521-29.